

Polyharmonic functions and random processes in cones

François Chapon

Université de Toulouse

Institut de Mathématiques de Toulouse, UMR CNRS 5219, 31062 Toulouse Cedex 9, France

francois.chapon@math.univ-toulouse.fr

Éric Fusy

CNRS

LIX, UMR CNRS 7161, École Polytechnique, 1 rue Honoré d'Estienne d'Orves, 91120 Palaiseau,

France

fusy@lix.polytechnique.fr

Kilian Raschel

CNRS

Institut Denis Poisson, UMR CNRS 7013, Université de Tours et Université d'Orléans, Parc de

Grandmont, 37200 Tours, France

raschel@math.cnrs.fr

Abstract

We investigate polyharmonic functions associated to Brownian motions and random walks in cones. These are functions which cancel some power of the usual Laplacian in the continuous setting and of the discrete Laplacian in the discrete setting. We show that polyharmonic functions naturally appear while considering asymptotic expansions of the heat kernel in the Brownian case and in lattice walk enumeration problems. We provide a method to construct general polyharmonic functions through Laplace transforms and generating functions in the continuous and discrete cases, respectively. This is done by using a functional equation approach.

2012 ACM Subject Classification Mathematics of computing → Markov processes; Theory of computation → Random walks and Markov chains; Mathematics of computing → Enumeration; Mathematics of computing → Generating functions

Keywords and phrases Brownian motion in cones; Heat kernel; Random walks in cones; Harmonic functions; Polyharmonic functions; Complete asymptotic expansions; Functional equations

Digital Object Identifier 10.4230/LIPIcs.AofA.2020.15

Funding *François Chapon*: F.C. is supported by the grant ANR-18-CE40-0006 MESA funded by the French National Research Agency (ANR).

Éric Fusy: É.F. is supported by the grant ANR-16-CE40-0009-01 GATO funded by the French National Research Agency (ANR).

Kilian Raschel: This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme under the Grant Agreement No. 759702.

Acknowledgements We would like to thank Cédric Lecouvey, Steve Melczer, Pierre Tarrago and Wolfgang Woess for very interesting discussions. This project has started in July 2019, when two authors were invited at the Institute of Mathematical Statistics of Münster University. The institute, and in particular Gerold Alsmeyer, is greatly acknowledged for hospitality. The first author also acknowledges the Institut Denis Poisson for the warm hospitality during his stay at the Université de Tours, where part of this work has been pursued. Finally, we thank the three anonymous referees for useful comments.



© François Chapon, Éric Fusy and Kilian Raschel;
licensed under Creative Commons License CC-BY

31st International Conference on Probabilistic, Combinatorial and Asymptotic Methods for the Analysis of Algorithms (AofA 2020).

Editors: Michael Drmota and Clemens Heuberger; Article No. 15; pp. 15:1–15:19



Leibniz International Proceedings in Informatics

Schloss Dagstuhl – Leibniz-Zentrum für Informatik, Dagstuhl Publishing, Germany

1 Introduction and motivations

In the continuous setting, polyharmonic functions are functions which cancel some power of the usual Laplacian. More precisely, a function v on some domain K of \mathbb{R}^d satisfying

$$\Delta^p v = 0$$

for some $p \geq 1$, where Δ is the usual Laplacian in \mathbb{R}^d , is said to be *polyharmonic* of order p , or *polyharmonic* for short. So polyharmonic functions of order 1 are just harmonic functions. Obviously, a polyharmonic function v_p of order p satisfies $\Delta v_p = v_{p-1}$, where v_{p-1} is polyharmonic of order $p-1$. For example, polynomials are polyharmonic. Harmonic functions have been tremendously investigated and pioneer works on polyharmonic functions go back to the work of Almansi [1]. One can consult for instance the monograph [2] for an introduction to this topic.

In particular, Almansi [1] proved that if the domain K is star-like with respect to the origin, then every polyharmonic function of order p admits a unique decomposition

$$f(x) = \sum_{k=0}^{p-1} |x|^{2k} h_k(x), \quad (1)$$

where each h_k is harmonic on K and $|x|$ is the Euclidean length of x , hence completely characterising continuous polyharmonic functions on such domains.

In comparison with the continuous case, much less is known in the discrete setting, where the Laplacian has to be replaced by a discrete difference operator. Some progress in understanding discrete polyharmonic functions has been made in the last two decades. For instance, one may cite [12], where the authors investigated polyharmonic functions for the Laplacian on trees, and proved a similar result as Almansi's theorem (1) for homogeneous trees. Recent works of Woess and co-authors [18, 21] are generalising this previous work.

Our original motivation to study discrete polyharmonic functions comes from the following framework. Consider a walk in \mathbb{Z}^d with step set \mathcal{S} confined in some cone $K \subset \mathbb{Z}^d$. Denote by $q(x, y; n)$ the number of n -length excursions between x and y staying in the cone K . To simplify, we only consider the case where y is the origin, but all considerations below can be generalised to $y \neq 0$. In various cases [15], the asymptotics of $q(x, 0; n)$ as $n \rightarrow \infty$ is known to admit the form

$$q(x, 0; n) \sim v_0(x) \gamma^n n^{-\alpha_0}, \quad (2)$$

where $v_0(x) > 0$ is a function depending only on x , $\gamma \in (0, |\mathcal{S}|]$ is the exponential growth, and α_0 is the critical exponent. It is easy to see that the function $v_0(x)$ in (2) defines a discrete harmonic function. Indeed, plugging (2) into the obvious recursive relation

$$q(x, 0; n+1) = \sum_{s \in \mathcal{S}} q(x+s, 0; n) \mathbf{1}_{\{x+s \in K\}}, \quad (3)$$

dividing by $\gamma^{n+1} n^{-\alpha_0}$ and letting $n \rightarrow \infty$, we obtain

$$v_0(x) = \frac{1}{\gamma} \sum_{s \in \mathcal{S}} v_0(x+s) \mathbf{1}_{\{x+s \in K\}}, \quad (4)$$

which proves that, with the assumption that $v_0(x) = 0$ for $x \notin K$, $v_0(x)$ is discrete harmonic for the Laplacian operator

$$Lf(x) = \frac{1}{\gamma} \sum_{s \in \mathcal{S}} f(x+s) - f(x), \quad (5)$$

that is, $Lv_0 = 0$. Denisov and Wachtel [15] go further and show that

- the exponential growth γ is $\min_{\mathbb{R}_+^d} \sum_{(s_1, \dots, s_d) \in \mathcal{S}} x_1^{s_1} \cdots x_d^{s_d}$, it does not depend on K ;
- the critical exponent α_0 equals $1 + \sqrt{\lambda_1 + (d/2 - 1)^2}$, where d is the dimension and λ_1 is the principal Dirichlet eigenvalue on some spherical domain constructed from K .

As a leading example, consider the simple random walk in the quarter plane, with step set $\{\leftarrow, \uparrow, \rightarrow, \downarrow\}$. In this case, the number of excursions $q((i, j), 0; n)$ is 0 if $m = \frac{n-i-j}{2}$ is not a non-negative integer, and otherwise takes the value

$$q((i, j), 0; n) = \frac{(i+1)(j+1)n!(n+2)!}{m!(m+i+j+2)!(m+i+1)!(m+j+1)!}, \quad (6)$$

see [9] and our Example 6. The equivalence (2) is then

$$q((i, j), 0; n) \sim \frac{4}{\pi} 4^n \frac{v_0(i, j)}{n^3}, \quad (7)$$

where $v_0(i, j) = (i+1)(j+1)$ is the well-known unique (up to multiplicative constants) harmonic function positive within the quarter plane with Dirichlet boundary conditions. Other examples of such asymptotics may be found for instance in [4, 10, 14].

Our aim in this discrete setting is to study more precise estimates than (2), by considering complete asymptotic expansions of the following form, as $n \rightarrow \infty$,

$$q(x, 0; n) \sim \gamma^n \sum_{p \geq 0} \frac{v_p(x)}{n^{\alpha_p}}. \quad (8)$$

From such an asymptotic expansion and using similar ideas as in (3), (4) and (5), it is rather easy to prove that the terms v_p are polyharmonic functions, in the sense that a power $L^k v_p$ of the Laplacian operator vanishes. We will provide examples of such asymptotic expansions (at least for the first terms) and of the set of exponents $\{\alpha_p\}_{p \geq 0}$ appearing in (8).

On the other hand, the functional equation approach has proved to be fruitful when studying random walk problems. The reference book on this topic is the monograph [16] by Fayolle, Iasnogorodski and Malyshev. This method has been used in [20] to construct harmonic functions, both in the discrete and continuous settings. Basically, the method consists of drawing from the harmonicity condition a functional equation satisfied by the generating function (in the discrete setting) or by the Laplace transform (in the continuous setting) of a harmonic function. Solving some boundary value problem for these quantities leads, via Cauchy or Laplace inversion, to the sought harmonic function. We will provide an implementation of this method to construct bi-harmonic functions, which can be generalised to polyharmonic functions.

The main features of our results are as follows:

- We shine a light on a new link between discrete polyharmonic functions and complete asymptotic expansions in the enumeration of walks.
- Our approach provides tools to study complete asymptotics expansions as in (8), but does not allow to prove their existence. On the other hand, the powerful approach of Denisov and Wachtel [15] seems restricted to the first term in the asymptotics (2). Indeed, one of the main tools in [15] is a coupling result of random walks by Brownian motion, which only provides an approximation of polynomial order, see [15, Lem. 17].
- We introduce a new class of functional equations (see (21) and (29)), for which the method of Tutte's invariants introduced in [23, 5, 6] proves to be useful.

■ In the unweighted planar case, it has been shown [8] that knowing the rationality of the exponent α_0 in (8) was sufficient to decide the non-D-finiteness of the series of excursions. However, for walks with big steps in dimension two or walk models in dimension three, this information is not enough [7]. As a potential application of our results, we might use arithmetic information on the other exponents α_p to study the algebraic nature, for example the transcendence, of the associated combinatorial series.

This paper is organised as follows. We choose to start with the continuous setting since computations are more enlightening and accessible. In Section 2, we prove that polyharmonic functions naturally arise when performing an asymptotic expansion of the Dirichlet heat kernel in a cone. We next present the functional equation method to construct polyharmonic functions. Our main result here is Theorem 4, where a class of solutions for the Laplace transform of a bi-harmonic function is provided. It shows that the Laplace transform of a bi-harmonic function can be expressed in terms of the Laplace transform of the related harmonic function plus some additional terms. This can be thought of as a Laplace transform version of Almansi's theorem (1). In Section 3, we exhibit the same phenomenon in the random walk setting. Discrete polyharmonic functions appear when considering the asymptotic expansion of coefficients counting walks with fixed endpoints in a domain, and the functional equation approach may be used to construct discrete polyharmonic functions.

These notes are the starting point of a long-term research project on discrete polyharmonic functions in cones. Notice that many ideas and techniques are not specific to cones and would work for many other domains of restriction K .

2 Classical polyharmonic functions and heat kernel expansions

As pointed out in [2, Chap. VI], the connection between the heat kernel and polyharmonic functions is very profound. Here, we deepen this connection by proving an exact asymptotic expansion for the heat kernel in terms of polyharmonic functions. We then implement the functional equation method to construct polyharmonic functions.

2.1 Exact asymptotic expansion for the Brownian semigroup in a cone

Let K be some cone in \mathbb{R}^d and consider the Brownian motion $(B_t)_{t \geq 0}$ killed at the boundary of K . Denote by $p(x, y; t)$ its transition density, that is the density probability function of the transition probability kernel

$$\mathbb{P}_x(B_t \in dy, \tau > t),$$

where τ is the first exit time of K . Recall the well-known fact that $p(x, y; t)$ corresponds to the heat kernel, i.e., the fundamental solution of the heat equation on K with Dirichlet boundary condition, see for instance [3]. Here, we prove that the heat kernel admits a complete asymptotic expansion in terms of polyharmonic functions for the Laplacian.

Denote by Δ the usual Laplacian on \mathbb{R}^d . In polar coordinates (r, θ) , where r is the radial part and θ the angular part, it writes:

$$\Delta = \frac{\partial^2}{\partial r^2} + \frac{d-1}{r} \frac{\partial}{\partial r} + \frac{1}{r^2} \Delta_{\mathbb{S}^{d-1}}, \quad (9)$$

where $\Delta_{\mathbb{S}^{d-1}}$ denotes the spherical Laplacian. Let respectively m_j and λ_j be the Dirichlet (normalised) eigenfunctions and eigenvalues for the spherical Laplacian on the generating set

163 $K \cap \mathbb{S}^{d-1}$, that is,

$$164 \quad \begin{cases} \Delta_{\mathbb{S}^{d-1}} m_j &= -\lambda_j m_j & \text{in } K \cap \mathbb{S}^{d-1}, \\ m_j &= 0 & \text{in } \partial(K \cap \mathbb{S}^{d-1}). \end{cases} \quad (10)$$

165 The eigenvalues satisfy $0 < \lambda_1 < \lambda_2 \leq \lambda_3 \leq \dots$ by [11, Chap. VII]. We introduce, for $j \geq 1$,

$$166 \quad \beta_j = \sqrt{\lambda_j + (d/2 - 1)^2} \quad \text{and} \quad b_j = 1 - d/2 + \sqrt{\lambda_j + (d/2 - 1)^2}. \quad (11)$$

167 Lemma 1 in [3] gives an explicit expression for the transition density $p(x, y; t)$ of the Brownian
168 motion in K . It states that, for $x, y \in \mathbb{R}^d$ and $t \in \mathbb{R}_+$,

$$169 \quad p(x, y; t) = \frac{\exp\left(-\frac{\rho^2 + r^2}{2t}\right)}{t(\rho r)^{\frac{d}{2}-1}} \sum_{j=1}^{\infty} I_{\beta_j}\left(\frac{\rho r}{t}\right) m_j(\theta) m_j(\eta), \quad (12)$$

170 where in polar coordinates $x = (\rho, \theta)$ and $y = (r, \eta)$. Here, I_β is the modified Bessel function
171 of the first kind of order β , satisfying the differential equation $I_\beta''(z) + \frac{1}{z}I_\beta'(z) = (1 + \frac{\beta^2}{z^2})I_\beta(z)$
172 and admitting the series expansion

$$173 \quad I_\beta(z) = \sum_{m=0}^{\infty} \frac{1}{m! \Gamma(m + \beta + 1)} \left(\frac{z}{2}\right)^{2m+\beta}. \quad (13)$$

174 The following easy lemma will allow us to define certain polyharmonic functions.

175 ► **Lemma 1.** *For any $\mu \geq 0$ and $j \geq 1$, let $f_{\mu,j}$ be defined in spherical coordinates by*

$$176 \quad f_{\mu,j}(r, \theta) = r^\mu m_j(\theta). \quad (14)$$

177 Then $f_{\mu,j}$ satisfies

$$178 \quad \Delta f_{\mu,j} = (\mu^2 + (d-2)\mu - \lambda_j) f_{\mu-2,j}. \quad (15)$$

179 **Proof.** The proof is elementary using (9) and (10). ◀

180 ► **Corollary 2.** *For any $k \in \mathbb{N}$, the function $f_{b_j+2k,j}$ defined in (14) is k -polyharmonic.*

181 **Proof.** It is obvious that $\mu = b_j$ satisfies $\mu^2 + (d-2)\mu - \lambda_j = 0$, see (11), so that $f_{b_j,j}$ is
182 harmonic by (15). An induction based on (15) completes the proof. ◀

183 Doing an expansion of the heat kernel (12) as $t \rightarrow \infty$ and using series expansions of the
184 exponential function and of the Bessel function (13), one immediately obtains:

185 ► **Theorem 3.** *The Dirichlet heat kernel $p(x, y; t)$ in K admits the following expansion, as
186 $t \rightarrow \infty$, where $f_{b_j+2k,j}$ is defined in (14), and b_j and β_j in (11):*

$$187 \quad p(x, y; t) \sim$$

$$188 \quad \sum_{j \geq 1} \sum_{k, m \geq 0} \sum_{n=0}^k \frac{1}{t^{1+\beta_j+k+2m}} \frac{(-1)^k \binom{k}{n}}{2^k k! m! \Gamma(m + \beta_j + 1)} f_{b_j+2(m+n),j}(\rho, \theta) f_{b_j+2(m+k-n),j}(r, \eta).$$

191 As such, the above result shows that the transition density of the Brownian motion
192 in K admits, as $t \rightarrow \infty$, an asymptotic expansion in descending powers of t and in terms

of polyharmonic functions for the Laplacian (see Corollary 2). Moreover, the set of these exponents is (with $\mathbb{N} = \{0, 1, 2, \dots\}$)

$$\bigcup_{j=1}^{\infty} (\beta_j + 1 + \mathbb{N}). \quad (16)$$

Note that, depending on the cone, there might be an overlap between the sets $\beta_j + 1 + \mathbb{N}$. For instance, in the quadrant in dimension 2, one has $\beta_j = 2j$ and the set in (16) reduces to $\{3, 4, 5, \dots\}$. On the other hand, in dimension 2 in a cone of opening α such that $\pi/\alpha \notin \mathbb{Q}$, there is no overlap between the points in (16).

As a last remark, we note that the same phenomenon appears for the survival probability $\mathbb{P}_x(\tau > t)$. Indeed, thanks to its explicit expression given by [3, Thm 1] (in terms of the confluent hypergeometric function), one can write down an asymptotic expansion of $\mathbb{P}_x(\tau > t)$ in descending powers of t in terms of polyharmonic functions for the Laplacian.

2.2 The functional equation approach

We apply here the functional equation approach in order to construct polyharmonic functions for the 2-dimensional killed Brownian motion in a convex cone. This approach has been previously introduced in [20] to compute harmonic functions, and is an adaptation of the functional equation method of the random walk case. Our main result is Theorem 4, which gives the general form of the Laplace transform of a bi-harmonic function.

Consider the Brownian motion B in the quarter plane \mathbb{R}_+^2 (compared to the last section, we use (x, y) for the coordinates of a 2d point) with covariance matrix

$$\Sigma = \begin{pmatrix} \sigma_{11} & \sigma_{12} \\ \sigma_{12} & \sigma_{22} \end{pmatrix},$$

with $\sigma_{11}, \sigma_{22} > 0$ and $\det \Sigma = \sigma_{11}\sigma_{22} - \sigma_{12}^2 \geq 0$. Its infinitesimal generator is the operator

$$\mathcal{G}f = \frac{1}{2} \left(\sigma_{11} \frac{\partial^2 f}{\partial x^2} + 2\sigma_{12} \frac{\partial^2 f}{\partial x \partial y} + \sigma_{22} \frac{\partial^2 f}{\partial y^2} \right).$$

Note that through some linear transformation ϕ (see [20, Eq. (5.1)]), one obtains the Brownian motion with identity covariance matrix in the cone $\phi(\mathbb{R}_+^2)$.

The *kernel* associated to the Brownian motion is defined as the quantity

$$\gamma(x, y) = \frac{1}{2}(\sigma_{11}x^2 + 2\sigma_{12}xy + \sigma_{22}y^2),$$

for $(x, y) \in \mathbb{C}^2$. The Laplace transform of a function f , which in the continuous case is the analogous quantity of the notion of generating function, is defined as

$$L(f)(x, y) = \iint_{[0, \infty)^2} f(u, v) e^{-(xu + yv)} du dv,$$

for $(x, y) \in \mathbb{C}^2$ with positive real parts.

Now, let h be a harmonic function associated with the Brownian motion with covariance matrix Σ , that is, h vanishes on the boundary axes of the quadrant and satisfies $\mathcal{G}h = 0$. The functional equation for h takes the following form (see [20, Eq. (A.1)]):

$$\gamma(x, y)L(h)(x, y) = \frac{1}{2}(\sigma_{11}L_1(h)(y) + \sigma_{22}L_2(h)(x)) + L(\mathcal{G}h)(x, y),$$

where we have denoted

$$\begin{cases} L_1(h)(y) &:= L\left(\frac{\partial h}{\partial x}(0, \cdot)\right)(y) = \int_0^\infty \frac{\partial h}{\partial x}(0, v) e^{-yv} dv, \\ L_2(h)(x) &:= L\left(\frac{\partial h}{\partial y}(\cdot, 0)\right)(x) = \int_0^\infty \frac{\partial h}{\partial y}(u, 0) e^{-xu} du. \end{cases}$$

Using the harmonicity condition $\mathcal{G}h = 0$, the functional equation for h rewrites as

$$\gamma(x, y)L(h)(x, y) = \frac{1}{2}(\sigma_{11}L_1(h)(y) + \sigma_{22}L_2(h)(x)). \quad (17)$$

We recall below the key argument of the method of [20] to solve the functional equation (17), which leads to harmonic functions for the Brownian motion via Laplace inversion. We will subsequently apply a related method to obtain polyharmonic functions.

Consider the two solutions of $\gamma(x, Y(x)) = 0$, which, since γ is a homogeneous polynomial of degree two, are explicitly given by $Y_\pm(x) = c_\pm x$, with

$$c_\pm = \frac{-\sigma_{12} \pm i\sqrt{\det \Sigma}}{\sigma_{22}}, \quad (18)$$

so that $c_+ = \overline{c_-}$. We write $c_\pm = ce^{\pm i\theta}$, with $c = \sqrt{\frac{\sigma_{11}}{\sigma_{22}}}$ and θ such that $\cos \theta = -\frac{\sigma_{12}}{\sqrt{\sigma_{11}\sigma_{22}}}$.

Denote by \mathcal{G}_Y the domain delimited by the curve $Y_+([0, \infty]) \cup Y_-([0, \infty]) = c_+[0, \infty] \cup c_-[0, \infty]$ and containing the positive axis $[0, \infty]$. Plugging each of the solutions $c_\pm x$ into the functional equation (17), one obtains a *boundary value problem* for $L_1(h)$, which states that:

1. $L_1(h)$ is analytic on \mathcal{G}_Y ,
2. $L_1(h)$ is continuous on $\overline{\mathcal{G}_Y} \setminus \{0\}$,
3. For all $x \in (0, \infty]$, $L_1(h)$ satisfies the boundary equation $L_1(h)(c_+x) = L_1(h)(c_-x)$.

In order to solve this problem, one introduces the conformal mapping ω from \mathcal{G}_Y onto $\mathbb{C} \setminus \mathbb{R}_-$ defined by $\omega(x) = x^{-\pi/\theta}$. One eventually obtains that a class of solutions is obtained by letting $L_1(h)$ to be of the form

$$L_1(h)(y) = P\left(\frac{1}{y^{\pi/\theta}}\right), \quad (19)$$

for any given polynomial P . The same applies to $L_2(h)$ (by considering the solutions of $\gamma(X(y), y) = 0$), and using the functional equation (17) and the fact that $(c_\pm)^{\pi/\theta} = -c^{\pi/\theta}$, one must have

$$L_2(h)(y) = -\frac{\sigma_{11}}{\sigma_{22}}P\left(-\frac{1}{c^{\pi/\theta}x^{\pi/\theta}}\right),$$

with the same P as in (19). Hence, using again the functional equation (17), we deduce that the Laplace transform of h writes

$$L(h)(x, y) = \frac{1}{2}\sigma_{11} \frac{P\left(\frac{1}{y^{\pi/\theta}}\right) - P\left(-\frac{1}{c^{\pi/\theta}x^{\pi/\theta}}\right)}{\gamma(x, y)}. \quad (20)$$

In particular, taking P to be a polynomial of degree 1, one gets

$$L(h)(x, y) = \frac{\sigma_{22} \frac{\mu_2}{x^{\pi/\theta}} + \sigma_{11} \frac{\mu_1}{y^{\pi/\theta}}}{\gamma(x, y)},$$

where the constants are related by $\mu_2 = \mu_1(\frac{\sigma_{22}}{\sigma_{11}})^{1-\pi/2\theta}$. Taking the inverse Laplace transform, one should recover the unique positive harmonic function (written in polar coordinates (ρ, η))

$$h(x, y) = \rho^{\frac{\pi}{\theta}} \sin\left(\frac{\pi}{\theta}\eta\right).$$

Suppose now that v is bi-harmonic and satisfies $\mathcal{G}v = h$, where h is harmonic. The functional equation for v now reads

$$\gamma(x, y)L(v)(x, y) = \frac{1}{2}(\sigma_{11}L_1(v)(y) + \sigma_{22}L_2(v)(x)) + L(h)(x, y). \quad (21)$$

By considering the roots of the kernel γ and using the same method as above, we obtain

$$\frac{1}{2}\sigma_{11}L_1(v)(c_+x) - \frac{1}{2}\sigma_{11}L_1(v)(c_-x) = L(h)(x, c_-x) - L(h)(x, c_+x). \quad (22)$$

We now have an *a priori* non-homogeneous boundary value problem for v , that we can in fact transform into an homogeneous one, thanks to the (already known) explicit form of $L(h)$. The key remark to this task is that $(c_+x)^{\pi/\theta} = (c_-x)^{\pi/\theta} = -(cx)^{\pi/\theta}$. Rewriting (20) as

$$L(h)(x, y) = \frac{\sigma_{11}}{\sigma_{22}} \frac{P\left(\frac{1}{y^{\pi/\theta}}\right) - P\left(\frac{1}{(c_{\pm}x)^{\pi/\theta}}\right)}{(y - c_-x)(y - c_+x)}$$

and letting $y \rightarrow c_+x$ and $y \rightarrow c_-x$, one finds

$$L(h)(x, c_{\pm}x) = \mp \frac{\sigma_{11}}{\sigma_{22}} \frac{\pi}{\theta} \frac{1}{(c_{\pm}x - c_{\mp}x)} P'\left(\frac{1}{(c_{\pm}x)^{\pi/\theta}}\right) \frac{1}{(c_{\pm}x)^{\pi/\theta+1}}.$$

Eventually, we get

$$\begin{aligned} & L(h)(x, c_-x) - L(h)(x, c_+x) \\ &= \frac{\sigma_{11}}{\sigma_{22}} \frac{\pi}{\theta} \left(\frac{1}{(c_+x - c_-x)} \frac{P'\left(\frac{1}{(c_+x)^{\pi/\theta}}\right)}{(c_+x)^{\pi/\theta+1}} - \frac{1}{(c_-x - c_+x)} \frac{P'\left(\frac{1}{(c_-x)^{\pi/\theta}}\right)}{(c_-x)^{\pi/\theta+1}} \right) \\ &= \frac{\sigma_{11}}{\sigma_{22}} \frac{\pi}{\theta} \left(\frac{c_+}{c_+ - c_-} P'\left(\frac{1}{(c_+x)^{\pi/\theta}}\right) \frac{1}{(c_+x)^{\pi/\theta+2}} - \frac{c_-}{c_- - c_+} P'\left(\frac{1}{(c_-x)^{\pi/\theta}}\right) \frac{1}{(c_-x)^{\pi/\theta+2}} \right) \\ &= -\frac{\sigma_{11}}{\sigma_{22}} \frac{\pi}{\theta} \frac{c_+c_-}{(c_+ - c_-)^2} \left(P'\left(\frac{1}{(c_+x)^{\pi/\theta}}\right) \frac{1}{(c_+x)^{\pi/\theta+2}} - P'\left(\frac{1}{(c_-x)^{\pi/\theta}}\right) \frac{1}{(c_-x)^{\pi/\theta+2}} \right), \end{aligned}$$

where the last equality follows from $(c_+x)^{\pi/\theta} = (c_-x)^{\pi/\theta}$. Therefore, the boundary value equation (22) is now homogeneous, and of the form

$$\frac{1}{2}\sigma_{11}L_1(v)(c_+x) - F(c_+x) = \frac{1}{2}\sigma_{11}L_1(v)(c_-x) - F(c_-x),$$

where F is equal on $Y_+([0, \infty]) \cup Y_-([0, \infty])$ to

$$F(y) = -\frac{\sigma_{11}}{\sigma_{22}} \frac{\pi}{\theta} \frac{c_+c_-}{(c_+ - c_-)^2} P'\left(\frac{1}{y^{\pi/\theta}}\right) \frac{1}{y^{\pi/\theta+2}}. \quad (23)$$

We note that the simpler case when $F(c_+x) = F(c_-x)$ occurs exactly when $c_+^2 = c_-^2$, i.e., θ is 0 or $\pi/2$. In this way, we obtain a boundary value problem analogous to the harmonic case, which, on the boundary of \mathcal{G}_Y except at 0, leads to

$$\frac{1}{2}\sigma_{11}L_1(v)(y) - F(y) = Q\left(\frac{1}{y^{\pi/\theta}}\right),$$

for any given polynomial Q . The same computation applies to $L_2(v)$. As such, using the equation (21), the Laplace transform of the bi-harmonic function v admits the following form:

290 ► **Theorem 4.** For any polynomials P and Q , the formula

$$291 \quad L(v)(x, y) = \frac{1}{\gamma(x, y)} \left[Q \left(\frac{1}{y^{\pi/\theta}} \right) - Q \left(\frac{1}{(c_+x)^{\pi/\theta}} \right) + G(x, y) + L(h)(x, y) \right]$$

292 is the Laplace transform $L(v)$ of a bi-harmonic function v satisfying $\mathcal{G}v = h$, where h is a
293 harmonic function with Dirichlet boundary conditions, where the Laplace transform $L(h)$ of
294 h has the form (20) and where

$$295 \quad G(x, y) = F(y) - F(c_+x) - L(h)(x, c_+x),$$

296 with F defined in Eq. (23).

297 The above theorem can be understood as a Laplace transform counterpart of Almansí's
298 theorem [1].

299 Recursively, if v_n is polyharmonic of order n with $\mathcal{G}v_n = v_{n-1}$, where v_{n-1} is polyharmonic
300 of order $n-1$, the above method permits to express the Laplace transform of v_n through
301 the one of v_{n-1} , allowing to construct polyharmonic functions via Laplace inversion.

302 Further computations for the Brownian motion with identity covariance matrix are
303 proposed in Appendix A.

304 **3 Discrete polyharmonic functions**

305 Similarly to the continuous setting, we first investigate the appearance of polyharmonic
306 functions in the asymptotic expansions of the counting coefficients of lattice paths with
307 prescribed endpoints, starting from an exact expression for these coefficients (such exact
308 expressions may typically be obtained from reflection principles). We then implement the
309 functional equation approach to construct polyharmonic functions.

310 Our framework is thus the following. We consider random walks in the quarter plane \mathbb{Z}_+^2
311 with the following assumptions:

- 312 1. The walk is homogeneous with transition probabilities $\{p_{i,j}\}_{-1 \leq i,j \leq 1}$ to the eight nearest
313 neighbours and $p_{0,0} = 0$ (so we are only considering walks with small steps),
- 314 2. In the list $p_{1,1}, p_{1,0}, p_{1,-1}, p_{0,-1}, p_{-1,-1}, p_{-1,0}, p_{-1,1}, p_{0,1}$, there are no three consecutive
315 zeros (to avoid degenerate cases),
- 316 3. The drifts $\sum_{i,j} i p_{i,j}$ and $\sum_{i,j} j p_{i,j}$ are zero.

317 The Markov operator P of the walk is defined on discrete functions by

$$318 \quad Pf(x, y) = \sum_{-1 \leq i,j \leq 1} p_{i,j} f(x+i, y+j),$$

319 and the Laplacian operator is $L = P - I$. A function f is said to be *harmonic* if $Lf = 0$ and
320 *polyharmonic* of order p if $L^p f = 0$.

321 **3.1 Examples of asymptotic expansion in walk enumeration problems**

322 We start by recalling a few exact expressions for the number of quarter plane walks of length
323 n with prescribed endpoints.

324 ► **Example 5** (The diagonal walk). The step set is $\{\nearrow, \nwarrow, \searrow, \swarrow\}$, with uniform transition
325 probabilities $\frac{1}{4}$. It is well known (see for instance [9]) that

$$326 \quad q((i, j), (0, 0); n) = \frac{(i+1)(j+1)}{\frac{n+i+2}{2} \frac{n+j+2}{2}} \binom{n}{\frac{n+i}{2}} \binom{n}{\frac{n+j}{2}}, \quad (24)$$

15:10 Polyharmonic functions in cones

with i and j having the same parity as n . Starting from (24), one can prove that

$$q((i, j), (0, 0); n) \sim \frac{8}{\pi} 4^n \sum_{p \geq 0} \frac{v_p(i, j)}{n^{3+p}}, \quad (25)$$

where the first few terms in the above asymptotic expansion are given by

$$\begin{cases} v_0(i, j) &= (i+1)(j+1), \\ v_1(i, j) &= -\frac{1}{2}(i+1)(j+1)(i^2 + j^2 + 2i + 2j + 9). \end{cases}$$

The first term v_0 is the well-known unique (up to multiplicative constants) positive harmonic function, with Dirichlet conditions; it is the same as for the simple walk, see (7) and (26). The next term satisfies $Lv_1 = -3v_0$, and therefore is bi-harmonic. Note that in fact, using the explicit expression of the Laplacian L , it is obvious that any polynomial of degree at most $2p-1$ is polyharmonic of order p , since for any polynomial f of degree k , Lf has degree at most $k-2$ (it is a discrete equivalent of Lemma 1).

To derive a full asymptotic expansion of (24), we shall use the Laplace method applied to the counting coefficients rewritten as an integral, in the spirit of [22, p. 75–79] (alternatively one can apply the saddle-point method [17, Chap. B VIII] in the framework of analytic combinatorics in several variables [14, 19]). We choose to postpone it to Appendix B, since the computations are a bit long, though straightforward.

► **Example 6** (The simple random walk). The step set is $\{\leftarrow, \uparrow, \rightarrow, \downarrow\}$, with uniform transition probabilities $\frac{1}{4}$. We have (6) by [9]. Again, starting from (6), one can prove that

$$q((i, j), (0, 0); n) \sim \frac{4}{\pi} 4^n \sum_{p \geq 0} \frac{v_p(i, j)}{n^{3+p}},$$

where the first few terms in the asymptotic expansion are

$$\begin{cases} v_0(i, j) &= (i+1)(j+1), \\ v_1(i, j) &= -\frac{1}{4}(i+1)(j+1)(2i^2 + 2j^2 + 4i + 4j + 15). \end{cases} \quad (26)$$

Again, v_0 is harmonic, and since $Lv_1 = -\frac{3}{2}v_0$, v_1 is bi-harmonic.

► **Example 7** (The tandem walk). The step set is $\{\nearrow, \rightarrow, \downarrow\}$ with uniform transition probabilities $\frac{1}{3}$. From [10, Prop. 9], we know that:

$$q((i, j), (0, 0); n) = \frac{(i+1)(j+1)(i+j+2)(3m+2i+j)!}{m!(m+i+1)!(m+i+j+2)!},$$

with $n = 3m + 2i + j$. In this case, writing the asymptotic expansion

$$q((i, j), (0, 0); n) \sim \frac{\sqrt{3}}{2\pi} 3^n \sum_{p \geq 0} \frac{v_p(i, j)}{n^{4+p}},$$

one has for the harmonic function v_0 and the bi-harmonic function v_1 ,

$$\begin{cases} v_0(i, j) &= (i+1)(j+1)(i+j+2), \\ v_1(i, j) &= -\frac{1}{9}(i+1)(j+1)(i+j+2)(3i^2 + 3j^2 + 3ij + 9i + 9j + 38). \end{cases} \quad (27)$$

3.2 Functional equation approach in the discrete case

We implement here the functional equation method to construct polyharmonic functions.

We start by recalling the key arguments in the harmonic case; details may be found in [20].

For a harmonic function h , we denote by H its generating function, namely,

$$H(x, y) = \sum_{i, j \geq 0} h(i, j) x^i y^j.$$

The *kernel* of the random walk is defined as the polynomial

$$K(x, y) = xy \left(\sum_{-1 \leq k, \ell \leq 1} p_{k, \ell} x^{-k} y^{-\ell} - 1 \right).$$

The harmonic equation $Lh = 0$ yields the following *functional equation*

$$K(x, y)H(x, y) = K(x, 0)H(x, 0) + K(0, y)H(0, y) - K(0, 0)H(0, 0). \quad (28)$$

To solve (28), one first proves that the function $H(x, 0)$ (and similarly $H(0, y)$) satisfies a *boundary value problem* (see [20]):

1. $H(x, 0)$ is analytic in \mathcal{G}_X ,
2. $H(x, 0)$ is continuous on $\overline{\mathcal{G}_X} \setminus \{1\}$,
3. For all x in the boundary of \mathcal{G}_X except at 1, $H(x, 0)$ satisfies the boundary equation:

$$K(x, 0)H(x, 0) - K(\bar{x}, 0)H(\bar{x}, 0) = 0.$$

Here, \mathcal{G}_X is a certain domain bounded by the curve $X_+([y_1, 1]) \cup X_-([y_1, 1])$, where $X_{\pm}(y)$ are the branches of the algebraic function defined by $K(X(y), y) = 0$. Indeed, writing K as

$$K(x, y) = \tilde{\alpha}(y)x^2 + \tilde{\beta}(y)x + \tilde{\gamma}(y),$$

where $\tilde{\alpha}, \tilde{\beta}, \tilde{\gamma}$ are polynomials of degree 2 whose coefficients depend on the model, we have

$$X_{\pm}(y) = \frac{-\tilde{\beta}(y) \pm \sqrt{\tilde{\delta}(y)}}{2\tilde{\alpha}(y)},$$

where $\tilde{\delta}(y) = \tilde{\beta}(y)^2 - 4\tilde{\alpha}(y)\tilde{\gamma}(y)$. The functions X_{\pm} are thus meromorphic on a cut plane, determined by the zeros of $\tilde{\delta}$.

It follows by [20] that $K(x, 0)H(x, 0)$ may be written as a function of a certain conformal mapping ω (see [20, Eq. (3.1)] for its explicit expression):

$$K(x, 0)H(x, 0) = P(\omega(x)),$$

where P is an arbitrary entire function, for example a polynomial. This represents the analogous statement as (19) in the continuous setting. By the functional equation (28), one eventually finds that

$$H(x, y) = \frac{P(\omega(x)) - P(\omega(X_+(x)))}{K(x, y)},$$

which again should be compared with (20) in the continuous case.

For a bi-harmonic function v , satisfying $Lv = h$ with h a harmonic function, the functional equation now writes

$$K(x, y)V(x, y) = K(x, 0)V(x, 0) + K(0, y)V(0, y) - K(0, 0)V(0, 0) - xyH(x, y), \quad (29)$$

15:12 Polyharmonic functions in cones

where V is the generating function of v , i.e., $V(x, y) = \sum_{i,j \geq 0} v(i, j)x^i y^j$; compare with (21). Notice that the equation (29) is very close to functional equations coming up in walk enumeration problems.

Plugging the roots of the kernel into (29), one has

$$K(X_{\pm}(y), 0)V(X_{\pm}(y), 0) + K(0, y)V(0, y) - K(0, 0)V(0, 0) - X_{\pm}(y)yH(X_{\pm}(y), y) = 0,$$

which leads to the boundary equation

$$K(x, 0)V(x, 0) - K(\bar{x}, 0)V(\bar{x}, 0) = y(xH(x, y) - \bar{x}H(\bar{x}, y)), \quad (30)$$

for x on the boundary of \mathcal{G}_X (except at 1).

Note that a general method to solve this kind of boundary value problem (30) exists [16], for any quantity in the right-hand side, ending up in some contour integral expression for the unknown function $K(x, 0)V(x, 0)$. We choose to provide below examples with simpler, integral-free expressions. Indeed, the resolution of (30) is made easier in some peculiar cases, for instance when the right-hand side of (30) is zero (which occurs for the simple random walk, see Example 6 below), or when it can be decoupled in the terminology of [6] (which is analogous to the continuous setting and holds for the tandem walk, see Appendix C).

► **Example 6 (continued).** We consider here the case of the simple random walk, with kernel

$$K(x, y) = xy \left(\frac{1}{4} \left(x + \frac{1}{x} + y + \frac{1}{y} \right) - 1 \right).$$

The domain \mathcal{G}_X is the open unit disk, and the conformal mapping ω admits the expression $\omega(x) = \frac{x}{(1-x)^2}$, see [20]. A computation shows that $\omega(X_+(y)) = -\omega(y)$, thus one gets that the generating function of a harmonic function h may be written as

$$H(x, y) = \frac{P(\omega(x)) - P(-\omega(y))}{K(x, y)}.$$

Choosing $P(x) = \frac{x}{4}$ leads to

$$H(x, y) = \frac{\frac{\frac{1}{4}x}{(1-x)^2} + \frac{\frac{1}{4}y}{(1-y)^2}}{xy \left(\frac{1}{4} \left(x + \frac{1}{x} + y + \frac{1}{y} \right) - 1 \right)} = \frac{1}{(1-x)^2(1-y)^2} = \sum_{i,j \geq 0} (i+1)(j+1)x^i y^j,$$

that is, H is the generating function of the unique positive harmonic function, see (26).

We now consider bi-harmonic functions. Using the explicit form of H , one sees that the right-hand side of Eq. (30) vanishes. Indeed, we have

$$\begin{aligned} & X_+(y)H(X_+(y), y) - X_-(y)H(X_-(y), y) \\ &= X_+(y) \frac{P'(\omega(X_+(y)))\omega'(X_+(y))}{\tilde{\alpha}(y)(X_+(y) - X_-(y))} - X_-(y) \frac{P'(\omega(X_-(y)))\omega'(X_-(y))}{\tilde{\alpha}(y)(X_-(y) - X_+(y))}, \end{aligned}$$

which is equal to zero since $\omega(X_+(y)) = \omega(X_-(y))$ and

$$X_+(y) \frac{\omega'(X_+(y))}{X_+(y) - X_-(y)} - X_-(y) \frac{\omega'(X_-(y))}{X_-(y) - X_+(y)} = 0$$

by straightforward computations. The boundary equation has thus exactly the same form as the one in the harmonic case, so we get that on the boundary of \mathcal{G}_X ,

$$K(x, 0)V(x, 0) = Q(\omega(x)),$$

for some polynomial Q . Using (twice) the functional equation (29), the general form for the generating function of a bi-harmonic v satisfying $Lv = h$, with h harmonic, is thus

$$V(x, y) = \frac{Q(\omega(x)) - Q(-\omega(y)) + X_+(y)yH(X_+(y), y) - xyH(x, y)}{K(x, y)},$$

with

$$H(x, y) = \frac{P(\omega(x)) - P(-\omega(y))}{K(x, y)} \quad \text{and} \quad H(X_+(y), y) = \frac{P'(\omega(X_+(y)))\omega'(X_+(y))}{\tilde{\alpha}(y)(X_+(y) - X_-(y))}.$$

For instance, taking $P(x) = x$ and Q the zero polynomial leads to the bi-harmonic function (non symmetrical in i and j)

$$v(i, j) = (i + 1)j(j + 1)(j + 2).$$

Indeed, one has

$$X_+(y)H(X_+(y), y) = -\frac{y}{(1 - y)^4},$$

so the generating function V writes

$$V(x, y) = \frac{-4y}{(1 - x)^2(1 - y)^4},$$

which is easily inverted. On the other hand, taking $P(x) = x$ and $Q(x) = -2x^2 - \frac{5}{2}x$, one obtains the bi-harmonic function

$$v(i, j) = (i + 1)(j + 1)(2i^2 + 2j^2 + 4i + 4j + 15),$$

which is (up to a multiplicative constant) the bi-harmonic function v_1 appearing in Eq. (26). Another example will be treated in Appendix C.

References

- 1 E. Almansi. Sull'integrazione dell'equazione differenziale $\Delta^{2n} = 0$. *Annali di Mat. (3)*, 2:1–51, 1899.
- 2 Nachman Aronszajn, Thomas M. Creese, and Leonard J. Lipkin. *Polyharmonic functions*. Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, New York, 1983. Notes taken by Eberhard Gerlach, Oxford Science Publications.
- 3 Rodrigo Bañuelos and Robert G. Smits. Brownian motion in cones. *Probab. Theory Related Fields*, 108(3):299–319, 1997. URL: <https://doi.org/10.1007/s0044000050111>, doi:10.1007/s0044000050111.
- 4 Cyril Banderier and Philippe Flajolet. Basic analytic combinatorics of directed lattice paths. *Theoretical Computer Science*, 281(1-2):37–80, 2002.
- 5 Olivier Bernardi and Mireille Bousquet-Mélou. Counting colored planar maps: algebraicity results. *J. Combin. Theory Ser. B*, 101(5):315–377, 2011. URL: <https://doi.org/10.1016/j.jctb.2011.02.003>, doi:10.1016/j.jctb.2011.02.003.
- 6 Olivier Bernardi, Mireille Bousquet-Mélou, and Kilian Raschel. Counting quadrant walks via Tutte's invariant method. *Preprint arXiv:1708.08215*, 2017.
- 7 Alin Bostan, Mireille Bousquet-Mélou, and Stephen Melczer. Counting walks with large steps in an orthant. *Preprint arXiv:1806.00968*, 2018.
- 8 Alin Bostan, Kilian Raschel, and Bruno Salvy. Non-D-finite excursions in the quarter plane. *J. Combin. Theory Ser. A*, 121:45–63, 2014. URL: <https://doi.org/10.1016/j.jcta.2013.09.005>, doi:10.1016/j.jcta.2013.09.005.

- 461 9 Mireille Bousquet-Mélou. Counting walks in the quarter plane. In *Mathematics and computer*
462 *science, II (Versailles, 2002)*, Trends Math., pages 49–67. Birkhäuser, Basel, 2002.
- 463 10 Mireille Bousquet-Mélou and Marni Mishna. Walks with small steps in the quarter plane.
464 In *Algorithmic probability and combinatorics*, volume 520 of *Contemp. Math.*, pages 1–39.
465 Amer. Math. Soc., Providence, RI, 2010. URL: <https://doi.org/10.1090/conm/520/10252>,
466 doi:10.1090/conm/520/10252.
- 467 11 Isaac Chavel. *Eigenvalues in Riemannian geometry*, volume 115 of *Pure and Applied Mathem-*
468 *atics*. Academic Press, Inc., Orlando, FL, 1984. Including a chapter by Burton Randol, With
469 an appendix by Jozef Dodziuk.
- 470 12 Joel M. Cohen, Flavia Colonna, Kohur Gowrisankaran, and David Singman. Polyharmonic
471 functions on trees. *Amer. J. Math.*, 124(5):999–1043, 2002. URL: [http://muse.jhu.edu/](http://muse.jhu.edu/journals/american_journal_of_mathematics/v124/124.5cohen.pdf)
472 [journals/american_journal_of_mathematics/v124/124.5cohen.pdf](http://muse.jhu.edu/journals/american_journal_of_mathematics/v124/124.5cohen.pdf).
- 473 13 Louis Comtet. *Advanced combinatorics. The art of finite and infinite expansions*. Dordrecht,
474 Holland - Boston, U.S.A.: D. Reidel Publishing Company. X, 343 p. Dfl. 65.00., 1974.
- 475 14 J. Courtiel, S. Melczer, M. Mishna, and K. Raschel. Weighted lattice walks and universality
476 classes. *J. Combin. Theory Ser. A*, 152:255–302, 2017. URL: [https://doi.org/10.1016/j.](https://doi.org/10.1016/j.jcta.2017.06.008)
477 [jcta.2017.06.008](https://doi.org/10.1016/j.jcta.2017.06.008), doi:10.1016/j.jcta.2017.06.008.
- 478 15 Denis Denisov and Vitali Wachtel. Random walks in cones. *Ann. Probab.*, 43(3):992–1044,
479 2015. URL: <https://doi.org/10.1214/13-AOP867>, doi:10.1214/13-AOP867.
- 480 16 Guy Fayolle, Roudolf Iasnogorodski, and Vadim Malyshev. *Random walks in the quarter*
481 *plane*, volume 40 of *Probability Theory and Stochastic Modelling*. Springer, Cham, second
482 edition, 2017. Algebraic methods, boundary value problems, applications to queueing systems
483 and analytic combinatorics. URL: <https://doi.org/10.1007/978-3-319-50930-3>, doi:10.
484 1007/978-3-319-50930-3.
- 485 17 Philippe Flajolet and Robert Sedgewick. *Analytic combinatorics*. Cambridge University
486 Press, Cambridge, 2009. URL: <https://doi.org/10.1017/CB09780511801655>, doi:10.1017/
487 CB09780511801655.
- 488 18 Thomas Hirschler and Wolfgang Woess. Polyharmonic functions for finite graphs and Markov
489 chains. *Preprint arXiv:1901.08376*, 2019. *Frontiers in Analysis and Probability: in the Spirit*
490 *of the Strasbourg-Zürich Meetings*, Springer (to appear).
- 491 19 Stephen Melczer and Mark C. Wilson. Higher dimensional lattice walks: connecting com-
492 binatorial and analytic behavior. *SIAM J. Discrete Math.*, 33(4):2140–2174, 2019. URL:
493 <https://doi.org/10.1137/18M1220856>, doi:10.1137/18M1220856.
- 494 20 Kilian Raschel. Random walks in the quarter plane, discrete harmonic functions and conformal
495 mappings. *Stochastic Process. Appl.*, 124(10):3147–3178, 2014. With an appendix by Sandro
496 Franceschi. URL: <https://doi.org/10.1016/j.spa.2014.04.013>, doi:10.1016/j.spa.2014.
497 04.013.
- 498 21 Ecaterina Sava-Huss and Wolfgang Woess. Boundary behaviour of λ -polyharmonic functions
499 on regular trees. *Preprint arXiv:1904.10290*, 2019.
- 500 22 F. Spitzer. *Principles of Random Walk*. 2nd ed, Springer, New York, 1976.
- 501 23 W. T. Tutte. Chromatic sums revisited. *Aequationes Math.*, 50(1-2):95–134, 1995. URL:
502 <https://doi.org/10.1007/BF01831115>, doi:10.1007/BF01831115.

A Detailed computations for the standard Brownian motion in the quadrant

Here we apply the functional equation approach to the case of the Brownian motion in the quarter plane with identity covariance matrix. The kernel γ is equal to $\gamma(x, y) = \frac{1}{2}(x^2 + y^2)$, so $c_{\pm} = \pm i$ and $\theta = \frac{\pi}{2}$, see (18). The functional equation (17) for h harmonic is then

$$(x^2 + y^2)L(h)(x, y) = L_1(h)(y) + L_2(h)(x),$$

which leads to

$$L(h)(x, y) = \frac{P\left(\frac{1}{y^2}\right) - P\left(-\frac{1}{x^2}\right)}{x^2 + y^2}.$$

In case when P is the degree 1 polynomial $P(x) = x$, one gets $L(h)(x, y) = \frac{1}{x^2 y^2}$ which is the Laplace transform of the well-known unique positive harmonic function within the quarter plane $h(x, y) = xy$.

More generally, the choice of $P(x) = -(2j)!(-x)^j$ leads to the Laplace transform (in Cartesian coordinates) of the harmonic function $f_{2j,j}$ defined in (14). Indeed, recall that $f_{2j,j}(\rho, \theta) = \rho^{2j} \sin(2j\theta)$, which is written in Cartesian coordinates as follows. Recall that the Chebyshev polynomial U_j of the second kind is defined as $U_j(\cos \theta) \sin \theta = \sin(j\theta)$, $j \geq 0$, and admits the expression

$$U_j(z) = z^j \sum_{k=0}^{\lfloor j/2 \rfloor} \binom{j+1}{2k+1} (1 - z^{-2})^k.$$

Hence, thanks to the explicit expression of U_{2j-1} , the harmonic function $f_{2j,j}$ can be written, in Cartesian coordinates $(x, y) = (\rho \cos \theta, \rho \sin \theta)$,

$$f_{2j,j}(x, y) = \sum_{k=0}^{j-1} (-1)^k \binom{2j}{2k+1} y^{2k+1} x^{2j-(2k+1)}.$$

The Laplace transform of $f_{2j,j}$ is now computed using $L(x^n y^k) = \frac{n!k!}{x^{n+1} y^{k+1}}$, and one obtains

$$L(f_{2j,j})(x, y) = (2j)! \sum_{k=0}^{j-1} (-1)^k \frac{1}{y^{2k+2} x^{2j-2k}} = (2j)! \frac{\left(\frac{1}{x^2}\right)^j - \left(-\frac{1}{y^2}\right)^j}{x^2 + y^2}. \quad (31)$$

For v bi-harmonic, the functional equation (21) is

$$(x^2 + y^2)L(v)(x, y) = L_1(v)(y) + L_2(v)(x) + 2L(h)(x, y),$$

and the general form of the Laplace transform of v writes

$$L(v)(x, y) = \frac{Q\left(\frac{1}{y^2}\right) - Q\left(-\frac{1}{x^2}\right) + \frac{2}{x^4} P'\left(-\frac{1}{x^2}\right) + 2 \frac{P\left(\frac{1}{y^2}\right) - P\left(-\frac{1}{x^2}\right)}{x^2 + y^2}}{x^2 + y^2}, \quad (32)$$

where P and Q are arbitrary polynomials. Choosing $P(x)$ equal to x and $Q(x)$ of degree 2, equal to x^2 , gives that

$$L(v)(x, y) = \frac{x^2 + y^2}{x^4 y^4} = \frac{1}{x^2 y^4} + \frac{1}{x^4 y^2},$$

which is the Laplace transform of the function $v(x, y) = (x^2 + y^2)xy$, which corresponds in polar coordinate (ρ, θ) to the bi-harmonic function $f_{4,2}(\rho, \theta) = \rho^4 \sin 2\theta$ defined in (14).

More generally, choosing

$$P(x) = (-1)^{j+1}(2j)!2(2j+1)x^j \quad \text{and} \quad Q(x) = (-1)^{j+1}(2j)!2(2j+1)jx^{j+1}$$

leads to the bi-harmonic function $f_{2j+2,j}$. Indeed, since $f_{2j+2,j}(x, y) = (x^2 + y^2)f_{2j,j}(x, y)$, one has, from the usual properties of the Laplace transform, that $L(f_{2j+2,j}) = \Delta L(f_{2j,j})$. As such, by applying the Laplacian to the Laplace transform of $f_{2j,j}$ given in (31), one obtains that

$$L(f_{2j+2,j})(x, y) = \frac{(2j)!2(2j+1)}{x^2y^2(x^2+y^2)^2} \left\{ (j+2)x^2y^2 \left(\left(\frac{1}{x^2} \right)^j - \left(\frac{-1}{y^2} \right)^j \right) + j \left(y^4 \left(\frac{1}{x^2} \right)^j - x^4 \left(\frac{-1}{y^2} \right)^j \right) \right\}.$$

Now, plugging the above choice of P and Q in Eq. (32) gives easily the formula.

B Complete asymptotic expansion for the diagonal walk

As an explicit example, we provide a complete asymptotic expansion for the number (24) of n -excursions from the origin to (i, j) for the diagonal walk with steps from $\{\nearrow, \nwarrow, \searrow, \swarrow\}$. A straightforward way to obtain such an asymptotic expansion is to apply the standard Laplace's method (see [17, p. 755]) using an integral representation of (24) (in [22, p. 75–79], this is applied to obtain first order asymptotic estimates in lattice paths enumeration problems). This leads to an explicit new family of polynomials $(v_p)_{p \geq 0}$ of increasing degree, where v_p is the polyharmonic function of order $p+1$ appearing in the expansion (25), see Corollary 9.

Let us first introduce the necessary notations. Projecting the walk onto the coordinate axes, one gets two decoupled prefixes of Dyck paths. Hence (24) is obtained by a simple application of the reflection principle in the one-dimensional case, which gives that the number of non-negative paths from 0 to λ with n steps is given by

$$m(\lambda, n) := \binom{n}{\frac{n+\lambda}{2}} - \binom{n}{\frac{n+\lambda+2}{2}} = \frac{\lambda+1}{\frac{n+\lambda+2}{2}} \binom{n}{\frac{n+\lambda}{2}}, \quad (33)$$

with $\lambda \equiv n \pmod{2}$. Using the simple integral representation of the binomial coefficient

$$\binom{n}{k} = \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{-ikt} (1 + e^{it})^n dt,$$

one readily obtains the following integral representation for $m(\lambda, n)$:

$$m(\lambda, n) = \frac{2}{\pi} \int_{-\pi/2}^{\pi/2} 2^n (\cos y)^n \sin((\lambda+1)y) \sin(y) dy. \quad (34)$$

Now define the sequence $(\alpha(m))_{m \geq 1}$ as

$$\alpha(m) = \frac{(4^m - 1)|B_{2m}|2^{2m}}{2m(2m)!}, \quad (35)$$

where the B_{2m} 's are the Bernoulli numbers, which can be defined through the Riemann zeta function at even integers:

$$\zeta(2m) = \frac{|B_{2m}|(2\pi)^{2m}}{2(2m)!}.$$

568 Define also, for $s \geq k \geq 0$,

$$569 \quad B_{s,k}^\alpha := B_{s,k}(\alpha(2), \dots, \alpha(s-k+2)), \quad (36)$$

570 the rational numbers obtained by evaluating the partial ordinary Bell polynomial in the
571 variables $\alpha(m+1)$. Recall that by definition, see for instance [13], the partial ordinary Bell
572 polynomials in the variables $(x_k)_{k \geq 1}$ are the polynomials obtained by performing the formal
573 double series expansion:

$$574 \quad \exp\left(u \sum_{m \geq 1} x_m t^m\right) = \sum_{n \geq k \geq 0} B_{n,k}(x_1, \dots, x_{n-k+1}) t^n \frac{u^k}{k!}.$$

575 Note that the polynomial $B_{n,k}$ contains $p(n,k)$ monomials, where $p(n,k)$ stands for the
576 number of partitions of n into k parts, see [13] for details and for an explicit expression of
577 these polynomials. Finally, define for $p \geq k \geq 0$,

$$578 \quad C_{k,p}^\alpha = \frac{1}{k!} \sum_{j=k}^p \frac{(-1)^j}{(2p-2j+1)!} B_{j,k}^\alpha. \quad (37)$$

579 We first give a complete asymptotic expansion for prefixes of Dyck paths.

580 ► **Theorem 8.** *Let $m(\lambda, n)$ be the number of non-negative paths from 0 to $\lambda \in \mathbb{Z}_+$ given by*
581 *(33). The following asymptotic expansion holds as $n \rightarrow \infty$:*

$$582 \quad m(\lambda, n) \sim 2\sqrt{2} \frac{2^n}{\sqrt{\pi}} \frac{1}{n^{3/2}} \sum_{j \geq 0} \frac{(-1)^j}{n^j} h_j(\lambda),$$

583 where for $j \geq 0$,

$$584 \quad h_j(\lambda) = \sum_{p=0}^j \sum_{k=0}^p \frac{(-1)^k}{(2(j-p)+1)!} C_{k,p}^\alpha m_{2(k+j+1)}(\lambda+1)^{2(j-p)+1}, \quad (38)$$

585 where $m_{2k} = \frac{(2k)!}{2^k k!}$ is the $2k$ -th Gaussian moment and $C_{k,p}^\alpha$ is defined in (37).

586 Hence, the above theorem gives, in the one-dimensional case, an asymptotic expansion of
587 the number of non-negative paths in terms of polyharmonic functions. Indeed, it is easily
588 seen that the polynomial h_j has degree $2j+1$, so is polyharmonic of order $j+1$ for the
589 one-dimensional Laplacian $Lf(x) = \frac{1}{2}(f(x+1) + f(x-1)) - f(x)$.

590 Since the number of n -excursions for the diagonal walk is the product of two numbers of
591 (decoupled) Dyck paths, one readily obtains the following corollary.

592 ► **Corollary 9.** *Let $q(0, (i, j); n)$ be the number of diagonal paths with n steps from the origin*
593 *to (i, j) and confined in the quadrant, given by (24). Then*

$$594 \quad q(0, (i, j); n) \sim \frac{8}{\pi} \frac{1}{n^3} 4^n \sum_{p \geq 0} \frac{(-1)^p}{n^p} v_p(i, j),$$

595 where, with h_k defined in (38),

$$596 \quad v_p(i, j) = \sum_{k=0}^p h_k(i) h_{p-k}(j).$$

Clearly, the polynomial function v_p has degree $2p + 1$ and thus is polyharmonic of order $p + 1$ for the Laplacian associated to the diagonal walk. The set of exponents (16) appearing in the asymptotic expansion is here $3 + \mathbb{N}$.

Proof of Theorem 8. To obtain the claimed asymptotic expansion, we apply the Laplace method as in [17, p. 755] to the integral representation of $m(\lambda, n)$ in (34). Indeed, the cosine function admits only one maximum in the interval $[-\frac{\pi}{2}, \frac{\pi}{2}]$, at $y = 0$, and the contribution to the integral outside any fixed segment containing 0 is exponentially small and as such can be discarded for an asymptotic consideration.

So, first, we perform the change of variable $\theta = \frac{y}{\sqrt{n}}$ to get

$$m(\lambda, n) = 2^n \frac{2}{\pi} \frac{1}{n^{1/2}} \int_{-\frac{\pi}{2}\sqrt{n}}^{\frac{\pi}{2}\sqrt{n}} \cos\left(\frac{y}{\sqrt{n}}\right)^n \sin\left(\frac{y}{\sqrt{n}}\right) \sin\left((\lambda + 1)\frac{y}{\sqrt{n}}\right) dy.$$

The next step is to consider an asymptotic expansion of the integrand as $n \rightarrow \infty$. Using the Weierstrass product formula for the cosine function,

$$\cos y = \prod_{k=1}^{\infty} \left(1 - \frac{4y^2}{\pi^2(2k-1)^2}\right)$$

and the Taylor series of the logarithm function, one has

$$\log \cos(y) = - \sum_{m \geq 1} \alpha(m) y^{2m},$$

where the sequence $(\alpha(m))_{m \geq 1}$ is defined in (35). Note that an interpretation of the sequence $(\alpha(m))_{m \geq 1}$ is that they correspond to the cumulant sequence of the Bernoulli distribution $\frac{1}{2}\delta_{+1} + \frac{1}{2}\delta_{-1}$. Now one has, using $\alpha(1) = \frac{1}{2}$ and the Taylor series of the exponential function,

$$\cos\left(\frac{y}{\sqrt{n}}\right)^n = \exp\left(n \log \cos\left(\frac{y}{\sqrt{n}}\right)\right) = e^{-y^2/2} \sum_{s \geq 0} \frac{1}{n^s} \sum_{k=0}^s \frac{(-1)^k}{k!} B_{s,k}^{\alpha} y^{2(k+s)},$$

where $B_{s,k}^{\alpha}$ is the partial ordinary Bell polynomial defined in (36). Now, using the Taylor series of the sine function, and after some elementary manipulations, one gets

$$\begin{aligned} \cos\left(\frac{y}{\sqrt{n}}\right)^n \sin\left(\frac{y}{\sqrt{n}}\right) \sin\left((\lambda + 1)\frac{y}{\sqrt{n}}\right) \\ = e^{-y^2/2} \frac{1}{n} \sum_{j \geq 0} \frac{(-1)^j}{n^j} \sum_{p=0}^j \sum_{k=0}^p (-1)^k \frac{C_{k,p}^{\alpha}}{(2(j-p)+1)!} y^{2(k+j)+2} (\lambda + 1)^{2(j-p)+1}, \end{aligned}$$

where $C_{k,p}^{\alpha}$ is defined in (37).

The next step in the Laplace method is to neglect the tails. Hence, we write

$$m(\lambda, n) \sim \frac{2}{\pi} \frac{2^n}{n^{3/2}} \sum_{j \geq 0} \frac{(-1)^j}{n^j} \sum_{p=0}^j \sum_{k=0}^p \frac{(-1)^k C_{k,p}^{\alpha}}{(2(j-p)+1)!} (\lambda + 1)^{2(j-p)+1} \int_{-\kappa_n}^{\kappa_n} e^{-y^2/2} y^{2(k+j)+2} dy,$$

where κ_n is chosen so that the error bounds are exponentially small (for instance one can choose arbitrarily $\kappa_n = n^{1/10}$). Completing the tails of the Gaussian integral, that is

$$\int_{-\kappa_n}^{\kappa_n} e^{-y^2/2} y^{2(k+j)+2} dy \sim \int_{\mathbb{R}} e^{-y^2/2} y^{2(k+j)+2} dy = \sqrt{2\pi} \frac{(2(k+j+1))!}{2^{k+j+1}(k+j+1)!} = \sqrt{2\pi} m_{2(k+j+1)},$$

where $m_{2k} = \frac{(2k)!}{2^k k!}$ is the $2k$ -th Gaussian moment, one finally obtains, with h_j defined in (38), that

$$m(\lambda, n) \sim 2\sqrt{2} \frac{2^n}{\sqrt{\pi}} \frac{1}{n^{3/2}} \sum_{j \geq 0} \frac{(-1)^j}{n^j} h_j(\lambda). \quad \blacktriangleleft$$

C The example of tandem walks

In this subsequent example, we consider the tandem walk with steps from $\{\nearrow, \rightarrow, \downarrow\}$, see Example 7. In this case, the functional equation approach admits a nicer form because the right-hand side of Eq. (30) can be decoupled, that is, can be written as $G(X_+(y)) - G(X_-(y))$, for some function G . The computations are close to the continuous case but are quite tedious. First, from [20], we know that the generating function H of a harmonic function h is of the form

$$H(x, y) = \frac{P(\omega(x)) - P(\omega(X_+(y)))}{K(x, y)}, \quad (39)$$

where the conformal mapping ω is given by $\omega(x) = \frac{x^2}{(1-x)^3}$. The unique positive harmonic function $v_0(i, j) = \frac{1}{2}(i+1)(j+1)(i+j+2)$ of (27) is obtained choosing $P(x) = \frac{1}{3}x$.

Using the general form of H , one has

$$\begin{aligned} yX_+(y)H(X_+(y), y) - yX_-(y)H(X_-(y), y) = \\ 3 \frac{yX_+(y)\omega'(X_+(y))}{X_+(y) - X_-(y)} P'(\omega(X_+(y))) - 3 \frac{yX_-(y)\omega'(X_-(y))}{X_-(y) - X_+(y)} P'(\omega(X_-(y))). \end{aligned}$$

Define now the *decoupling* function on \mathcal{G}_X :

$$F(x) = -\frac{\omega(x)^2}{x} = -\frac{x^3}{(1-x)^6}. \quad (40)$$

Some computations show that

$$\frac{yX_+(y)\omega'(X_+(y))}{X_+(y) - X_-(y)} - \frac{yX_-(y)\omega'(X_-(y))}{X_-(y) - X_+(y)} = F(X_+(y)) - F(X_-(y)).$$

Minding the fundamental fact that $\omega(X_+(y)) = \omega(X_-(y))$, it follows that

$$yX_+(y)H(X_+(y), y) - yX_-(y)H(X_-(y), y) = G(X_+(y)) - G(X_-(y)),$$

where $G(x) = 3F(x)P'(\omega(x))$. One deduces that the generating function $V(x, y)$ for a bi-harmonic function v satisfying $Lv = h$ admits the form

$$\frac{1}{K(x, y)} \left(Q(\omega(x)) - Q(\omega(X_+(y))) + G(x) - G(X_+(y)) + X_+(y)yH(X_+(y), y) - xyH(x, y) \right),$$

where H has the general form given by Eq. (39) and $G(x) = 3F(x)P'(\omega(x))$ with the decoupling function F defined in Eq. (40). Note that this has to be compared with Theorem 4.

Choosing $P(x) = x$ and $Q = 0$ leads to the bi-harmonic function

$$v(i, j) = (j+1)(i+1)(i+j+2)(2i^3 + 3i^2j + 14i^2 + 5ij + 24i - 3ij^2 - 2j^3 - 4j^2 + 6j). \quad (41)$$

To obtain to bi-harmonic function v_1 of (27), one chooses $P(x) = -\frac{8}{9}x$ and $Q(x) = \frac{8}{3}x^2 + \frac{76}{27}x$. This is obtained by noticing that an appropriate linear combination of the bi-harmonic function (41) and of v_1 is harmonic and its generating function corresponds to the term

$$\frac{Q(\omega(x)) - Q(\omega(X_+(y)))}{K(x, y)}.$$

As such, computing its generating function leads to the polynomial Q .