


Bijjective Links on Planar Maps via Orientations

Éric Fusy


Dept. Math, University of British Columbia

Planar maps

- **Planar map** = graph drawn in the plane **without edge-crossing**, taken **up to isotopy** (continuous structure-preserving transformation)




- **Rooted map** = map + root edge




- **Motivations**: mesh compression, graph drawing + nice combinatorial properties


Families of planar maps




Planar map




Loopless map



Nonseparable map
(no separating vertex)




Triangulation




Irreducible triangulation
(no separating triangle) . – p.3/26

Families of planar maps




Planar map


separating
vertex




Loopless map



Nonseparable map
(no separating vertex)




Triangulation




Irreducible triangulation
(no separating triangle) · - p.3/26

Families of planar maps




Planar map


separating
vertex



Loopless map




Nonseparable map
(no separating vertex)



Triangulation


separating
triangle



Irreducible triangulation
(no separating triangle)


Enumeration of planar maps

- **Symbolic approach:** Tutte, Brown
- **Bijjective approach:** Cori, Schaeffer, Bouttier-Di Francesco-Guitter




Planar maps

$$\#(n \text{ edges}) = \frac{2 \cdot 3^n (2n)!}{(n+2)! n!}$$




Eulerian

$$\#(n \text{ edges}) = \frac{3 \cdot 2^{n-1} (2n)!}{(n+2)! n!}$$




Loopless

$$\#(n \text{ edges}) = \frac{2(4n+1)!}{(n+1)!(3n+2)!}$$




Nonseparable

$$\#(n \text{ edges}) = \frac{4(3n-3)!}{(n-1)!(2n)!}$$




4-regular

$$\#(n \text{ vert.}) = \frac{2 \cdot 3^n (2n)!}{(n+2)! n!}$$




Bicubic

$$\#(2n \text{ vert.}) = \frac{3 \cdot 2^{n-1} (2n)!}{(n+2)! n!}$$



Triangulations

$$\#(n+3 \text{ vert.}) = \frac{2(4n+1)!}{(n+1)!(3n+2)!}$$




Irreducible

$$\#(n+3 \text{ vert.}) = \frac{4(3n-3)!}{(n-1)!(2n)!}$$


Enumeration of planar maps

- **Symbolic approach:** Tutte, Brown
- **Bijective approach:** Cori, Schaeffer, Bouttier-Di Francesco-Guitter




Planar maps

$$\#(n \text{ edges}) = \frac{2 \cdot 3^n (2n)!}{(n+2)! n!}$$




Eulerian

$$\#(n \text{ edges}) = \frac{3 \cdot 2^{n-1} (2n)!}{(n+2)! n!}$$




Loopless

$$\#(n \text{ edges}) = \frac{2(4n+1)!}{(n+1)!(3n+2)!}$$




Nonseparable

$$\#(n \text{ edges}) = \frac{4(3n-3)!}{(n-1)!(2n)!}$$




4-regular

$$\#(n \text{ vert.}) = \frac{2 \cdot 3^n (2n)!}{(n+2)! n!}$$




Bicubic

$$\#(2n \text{ vert.}) = \frac{3 \cdot 2^{n-1} (2n)!}{(n+2)! n!}$$



Triangulations

$$\#(n+3 \text{ vert.}) = \frac{2(4n+1)!}{(n+1)!(3n+2)!}$$



Irreducible

$$\#(n+3 \text{ vert.}) = \frac{4(3n-3)!}{(n-1)!(2n)!}$$

①


②

③

④

Enumeration of planar maps

- **Symbolic approach:** Tutte, Brown
- **Bijective approach:** Cori, Schaeffer, Bouttier-Di Francesco-Guitter




① , ② well known bijections (Tutte) ③ recursive bijection (Wormald)

This talk:


- **new** bijective construction for ③
- **first** bijective construction for ④

Well known bijections

① Radial mapping (Tutte)




② Trinity mapping (Tutte)




Overview of the talk



- 1) Bijection **nonseparable maps** $\simeq$ **irreducible triang**
+ new duality relation for bipolar orientations




- 2) Bijection **loopless maps** $\simeq$ **triangulations**



nonseparable components



irreducible components




- 3) Applications to random generation and encoding

Bijection between nonseparable maps and irreducible triangulations


Bipolar orientations

Bipolar orientation = acyclic orientation with a unique source and a unique sink



Bipolar orientations


Bipolar orientation = acyclic orientation with a unique source and a unique sink



A map admits a bipolar orientation iff there is no separating vertex (nonseparable)


Transversal structures

Transversal structure = partition of inner edges into a red and a blue bipolar orientations that are transversal (introduced by Xin He'93)



Transversal structures



Transversal structure = partition of inner edges into a red and a blue bipolar orientations that are transversal (introduced by Xin He'93)




A triangulation of the 4-gon admits a transversal structure iff there is no separating triangle (irreducible)

Reformulating the bijection

Nonseparable




Irreducible



Reformulating the bijection



Nonseparable



Ossona de
Mendez'94




No pattern



Bipolar orientations



Irreducible




Fusy'05



No patterns




Transversal structures





Reformulating the bijection


Nonseparable




Ossona de
Mendez'94





Bipolar orientations



Irreducible




Fusy'05




Transversal structures



How the bijection works




How the bijection works




Bipolar $\rightarrow$ Bipolar intransitive

Start with a plane bipolar orientation




Bipolar $\rightarrow$ Bipolar intransitive

Double the root edge




Bipolar $\rightarrow$ Bipolar intransitive

Insert a white vertex in each edge




Bipolar $\rightarrow$ Bipolar intransitive


Triangulate the faces by **red edges**



Bipolar $\rightarrow$ Bipolar intransitive




Bipolar $\rightarrow$ Bipolar intransitive




Bipolar $\rightarrow$ Bipolar intransitive

A first bijection:




- 1) place a black vertex in each face
- 2) insert one black edge for each white vertex




Bipolar $\rightarrow$ Bipolar intransitive

A first bijection:



- 1) place a black vertex in each face
- 2) insert one black edge for each white vertex




Remark: These are counted by the **Baxter number**:

$$B_n = \frac{2}{n(n+1)^2} \sum_{k=0}^{n-1} \binom{n+1}{k} \binom{n+1}{k+1} \binom{n+1}{k+2}$$


Bipolar intransitive $\rightarrow$ Trans. struct.

Start with an **intransitive** bipolar orientation




Bipolar intransitive $\rightarrow$ Trans. struct.

Triangulate the faces by blue edges





Bipolar intransitive $\rightarrow$ Trans. struct.






Bipolar intransitive $\rightarrow$ Trans. struct.


A second bijection:




The bijection




The bijection



The bijection




The bijection



Bijection between loopless maps and triangulations


Decomposing a loopless map

- Block decomposition




Decomposing a loopless map

- Block decomposition




- For rooted loopless maps:



Nonseparable core
where each corner
is possibly occupied
by a loopless map


Decomposing a triangulation

- Classical decomposition at separating triangles



Decomposing a triangulation


- Classical decomposition at separating triangles



4-connected triangulation
where each face is
possibly occupied
by a triangulation


Decomposing a triangulation

- Classical decomposition at separating triangles




4-connected triangulation
where each face is
possibly occupied
by a triangulation

- **Here:** the same **after deleting an outer edge**




Decomposing a triangulation

- Classical decomposition at separating triangles



4-connected triangulation
where each face is
possibly occupied
by a triangulation

- Here: the same after deleting an outer edge



Irreducible triangulation
where each face is
possibly occupied
by a triangulation


The decompositions are parallel

A **loopless map** M ($|M| = \# \text{ edges}$) is

i) the vertex-map $\bullet$


or

ii) $|M| \geq 1$



=

Nonseparable core C



+

$2|C|$ loopless maps $M_1, \dots, M_{2|C|}$




$$|M| = |C| + \sum_{i=1}^{2|C|} |M_i|$$

A **triangulation** T ($|T| = \# \text{ vert.} - 3$) is

i) the triangle-map 


or

ii) $|T| \geq 1$



=

Irreducible core I



+

$2|I|$ triangulations $T_1, \dots, T_{2|I|}$



$$|T| = |I| + \sum_{i=1}^{2|I|} |T_i|$$


The decompositions are parallel

A **loopless map** M ($|M| = \# \text{ edges}$) is

i) the vertex-map $\bullet$


or

ii) $|M| \geq 1$



=

Nonseparable core C



+

$2|C|$ loopless maps $M_1, \dots, M_{2|C|}$




$$|M| = |C| + \sum_{i=1}^{2|C|} |M_i|$$

A **triangulation** T ($|T| = \# \text{ vert.} - 3$) is

i) the triangle-map


or

ii) $|T| \geq 1$



=

Irreducible core I



+

$2|I|$ triangulations $T_1, \dots, T_{2|I|}$



$$|T| = |I| + \sum_{i=1}^{2|I|} |T_i|$$

Nonseparable maps n edges

$\longleftrightarrow$ *bijection*


Irreducible triang. $n + 3$ vertices

Loopless maps n edges


$\longleftrightarrow$ *bijection*

Triangulations $n + 3$ vertices

Results obtained so far




Results obtained so far



Results obtained so far


rooted nonseparable
maps with n edges

*bipolar
orientations*




rooted irreducible
triangulations
with $n+3$ vertices

$$|\mathcal{N}_n| = |\mathcal{I}_n| = ?$$



rooted loopless maps
with n edges

*parallel
decompositions*




rooted triangulations
with $n+3$ vertices

$$|\mathcal{L}_n| = |\mathcal{T}_n| = ?$$


Counting the families

Irreducible triang. $\leftrightarrow$ ternary trees




**Fusy'05: irreducible triangulations
are in bijection with ternary trees**

Irreducible triang. $\leftrightarrow$ ternary trees




canonical
transversal structure
→




Fusy'05: irreducible triangulations
are in bijection with ternary trees



Irreducible triang. $\leftrightarrow$ ternary trees




canonical
transversal structure




Fusy'05: **irreducible triangulations**
are in bijection with ternary trees





Irreducible triang. $\leftrightarrow$ ternary trees




canonical
transversal structure




Fusy'05: irreducible triangulations
are in bijection with ternary trees



rooting




Triangulations $\leftrightarrow$ quaternary trees




Poulalhon-Schaeffer'03, Fusy-Poulalhon-Schaeffer'05:
triangulations are in bijection with quaternary trees

Triangulations $\leftrightarrow$ quaternary trees




Poulalhon-Schaeffer'03, Fusy-Poulalhon-Schaeffer'05:
triangulations are in bijection with quaternary trees

Triangulations $\leftrightarrow$ quaternary trees




Poulalhon-Schaeffer'03, Fusy-Poulalhon-Schaeffer'05:
triangulations are in bijection with quaternary trees

Triangulations $\leftrightarrow$ quaternary trees




Poulalhon-Schaeffer'03, Fusy-Poulalhon-Schaeffer'05:
triangulations are in bijection with quaternary trees

Triangulations $\leftrightarrow$ quaternary trees




Poulalhon-Schaeffer'03, Fusy-Poulalhon-Schaeffer'05:
triangulations are in bijection with quaternary trees


Triangulations $\leftrightarrow$ quaternary trees



Poulalhon-Schaeffer'03, Fusy-Poulalhon-Schaeffer'05:
 triangulations are in bijection with quaternary trees



Enumerative Results



Enumerative Results

rooted nonseparable
maps with n edges



rooted irreducible
triangulations
with $n+3$ vertices



rooted ternary
trees with $n-1$ nodes



$$|\mathcal{N}_n| = |\mathcal{T}_n| = \frac{4(3n-3)!}{(n-1)!(2n)!}$$



rooted loopless maps
with n edges



rooted triangulations
with $n+3$ vertices




rooted quaternary
trees with n nodes



$$|\mathcal{L}_n| = |\mathcal{I}_n| = \frac{2(4n+1)!}{(n+1)!(3n+2)!}$$


Enumerative Results



Applications


Algorithmic applications

- **General scheme:** (Schaeffer'99, Poulalhon-Schaeffer'03)




Algorithmic applications

- **General scheme:** (Schaeffer'99, Poulalhon-Schaeffer'03)




Algorithmic applications

- **General scheme:** (Schaeffer'99, Poulalhon-Schaeffer'03)



Algorithmic applications


- **General scheme:** (Schaeffer'99, Poulalhon-Schaeffer'03)



- Applies here to :
 - irreducible triangulations
 - triangulations

Algorithmic applications

- **General scheme:** (Schaeffer'99, Poulalhon-Schaeffer'03)



- Applies here to :
 - irreducible triangulations
 - triangulationsas well as :
 - loopless maps
 - nonseparable maps
- Curved arrows point from "irreducible triangulations" to "loopless maps" and from "nonseparable maps" to "irreducible triangulations".